


Upper Rim Allyl- and Arylazo-Coupled Calix[4]arenes as Highly Sensitive Chromogenic Sensors for Hg²⁺ Ion

Tsui-Lien Kao,[†] Chiung-Chiu Wang,[†] Yu-Ting Pan,^{†,‡} Ya-Jiun Shiao,[†] Jhy-Yuan Yen,[†] Chun-Mei Shu,[†] Gene-Hsiang Lee,[§] Shie-Ming Peng,[§] and Wen-Sheng Chung^{*,†,‡}

Department of Applied Chemistry, Center for Interdisciplinary Molecular Science, National Chiao Tung University, Hsinchu, Taiwan 30050, R.O.C., and Department of Chemistry, National Taiwan University, Taipei, Taiwan 106, R.O.C.

wschung@cc.nctu.edu.tw

Received November 30, 2004



The syntheses and chromogenic properties of calix[4]arenes, carrying 5,17-bisallyl-11,23-bis(*p*-X-phenyl)azo **3a–c**, 5,11,17-triallyl-23-(*p*-X-phenyl)azo **4a–c**, and 5,17-bis(hydroxypropyl)-11,23-bis(*p*-X-phenyl)azo groups on the upper rims **5a,b**, are described. Unexpectedly, UV/vis spectra of the very popular 4-(4-nitrophenyl)azophenol-coupled calix[4]arenes **3c** and **4c** did not show any shift in λ_{max} when 10 different metal perchlorates were added separately to the host in a methanol–chloroform ($v/v = 1/399$) cosolvent. In contrast, the absorption spectra of calix[4]arenes with either 4-methoxyphenylazo (**3b–5b**) or 4-phenylazo (**3a–5a**) on the upper rim showed substantial bathochromic shifts ($\Delta\lambda = 128–162$ nm) upon the addition of soft metal ions (such as Hg²⁺, Cr³⁺, and Cu²⁺). The 4-(4-methoxyphenyl)azophenol-coupled calix[4]arenes (the **3b–5b** series) are found to be highly sensitive for mercury ion (Hg²⁺) among the 10 different metal ions screened. Strong interactions between Hg²⁺ ion and the 4-(4-methoxyphenyl)azophenol(s) as well as the *p*-allyl groups are corroborated by the ¹H NMR studies of **3a,b**·Hg²⁺ complexes. Furthermore, Job's plots revealed 1:1 binding stoichiometry for all these *p*-allyl- and arylazo-coupled calix[4]arenes with transition metal ions, and Benesi–Hilderbrand plots were used for the determination of their association constants.

Introduction

The specific color change of a ligand upon metal complexation is an informative signal that can be utilized in an ion-sensing system.¹ Vögtle and Takagi independently pioneered the use of 4-(4-nitrophenyl)azo-coupled crowns and azacrowns as chromoionophores, which showed large UV/vis band shifts when cations were added.^{1a,b} In their work, the construction principle was as follows: mono- and diprotic chromophores were set into a crown

skeleton in such a way that the complexation of positively charged metal ions was accompanied by the dissociation of protons of the chromophore. The crown ether concept has been elegantly extended into supramolecular chemistry by the development of more rigid and selective systems such as cryptand,^{1c} spherand,^{1d} hemispherand,^{1e} and calixarene derivatives.^{1f–i,2}

Most of the research on calix[4]arenes as chromogenic ionophores involves the modification of the lower rim with metal-chelating groups such as acids, amides, esters, ketones, or any ligand that contains lone pair electrons, and then a fluorophore or chromophore is attached on the upper rim.^{1–4} Shinkai and co-workers were the first to synthesize calix[4]arene with a 4-(4-nitrophenyl)-

[†] Department of Applied Chemistry, National Chiao Tung University.

[‡] Center for Interdisciplinary Molecular Science, National Chiao Tung University.

[§] National Taiwan University.

azophenol unit and three ether ester residues on the lower rim. They found that on cation complexation in the presence of imidazole, the complex exhibits a new absorption maximum at 600 nm, which is a “perfect” lithium sensor.^{2a} It should be noted that imidazole alone does not cause any spectral change, confirming that deprotonation and complexation are integral events in the chromogenic response. A gas sensor for colorimetric determination of trimethylamine was later devised by McCarrick et al.^{2b} using a calix[4]arene bearing 4-(4-nitrophenyl)azophenol residues on the lower rim. Furthermore, Reinhoudt et al. also reported that a calix[4]arene with monoalkylated 4-(4-nitrophenyl)azophenol and triamides on the lower rim is a highly selective Pb²⁺ sensor, in which the direction of the shift was dependent on the conformation of the calixarenes.⁴

Undoubtedly, 4-(4-nitrophenyl)azophenol is one of the most popular and thoroughly studied chromogenic centers in chromoionophores.^{2,4} However, the azo group is usually considered to be only a chromogenic center, not a metal-chelating site; therefore, lower rim modification of calix[4]arene is usually required to endow it with metal-chelating ability (vide supra).²⁻⁴ Herein, we describe the synthesis and spectroscopic evaluation of eight upper rim allyl- and arylazo-coupled calix[4]arenes **3a–c**, **4a–c**, and **5a,b**, without lower rim modification. Surprisingly, those with 4-phenyl- or 4-(4-methoxyphenyl)-azophenol-coupled calix[4]arenes (**3a–5a** and **3b–5b** series, respectively) were found to be excellent chromogenic ionophores for transition metal ions, especially for Hg²⁺ ion. The Hg²⁺ detection by **3b–5b** gives rise to

marked changes in the absorption spectra (from light yellow to bright red), which are clearly visible to the naked eye. However, for 4-(4-nitrophenyl)azophenol-coupled calix[4]arenes (the **3c–4c**), no transition-metal ion sensing was observed. Lower rim modification of **3b** and **3c** with ether-esters gave chromoionophores **6b** and **6c**, which were also used in metal ion screening.

Results and Discussion

We have followed literature procedures for the preparation of compounds **1**,^{5a,b} **2**,^{5b} and **3a–c**.^{5c,d} Besides traditional organic spectroscopic identification (¹H and ¹³C NMR, MS, and HRMS spectra) of all these calix[4]arenes, single-crystal X-ray analysis of **3a** and **3c** confirmed their structures to be in cone conformations (Figure 1). Calix[4]arenes **4a–c** were synthesized in 73–91% yields using a method similar to that used in the preparation of **3a–c** (Scheme 1). Hydroboration⁶ of the allyl groups on **3a** and **3b** using 4 equiv of BH₃·THF followed by hydrogen peroxide treatment under basic conditions gave **5a** and **5b** in 39 and 58% yields, respectively. Furthermore, lower rim esterification⁷ of azo **3b** and **3c** by sodium hydride and ethyl bromoacetate in large excess (20 equiv) gave **6b** and **6c** in 74 and 36% yields, respectively (Scheme 2). The presence of the doublets (or broad singlets) around δ 4.2–4.4 and 3.4–3.7 in the ¹H NMR spectra^{1f} and two methylene bridge resonances (only one for compounds **6b,c**) at about δ 3.2 in the ¹³C NMR spectra^{1f} for compounds **3a–c**, **4a–c**, **5a,b**, and **6b,c** indicated that they are in cone conformations (see Experimental Section for details).

It is known that 4-(*p*-X)-phenylazo-1-naphthol undergoes an azo/hydrazone tautomerization in polar media, and the quinone-hydrazone form is found to be highly favored in polar solvents. Moreover, the azo/hydrazone tautomerism is found to be more favored in *para*-nitrophenyl azophenol but is least favored in *para*-methoxyphenyl azophenol (vide infra).^{8,9} To understand the tautomerism of these chromogenic azocalix[4]arenes

(1) (a) Löhr, H. G.; Vögtle, F. *Acc. Chem. Res.* **1985**, *18*, 65 and references therein. (b) Hayashita, T.; Takagi, M. In *Comprehensive Supramolecular Chemistry*; Atwood, J. L.; Davies, J. E. D., MacNicol, D. D., Vögtle, F., Suslick, K. S., Eds.; Pergamon: Oxford, 1996; pp 635–669. (c) Zazulak, W.; Chapoteau, E.; Czech, B. P.; Kumar, A. *J. Org. Chem.* **1992**, *57*, 6720. (d) Kaneda, T.; Umeda, S.; Tanigawa, H.; Misumi, S.; Kai, Y.; Morii, H.; Miki, K.; Kasai, N. *J. Am. Chem. Soc.* **1985**, *107*, 4802. (e) Helgeson, R. C.; Czech, B. P.; Chapoteau, E.; Gebauer, C. R.; Kumar, A.; Cram, D. J. *J. Am. Chem. Soc.* **1989**, *111*, 6339. (f) Gutsche, C. D. *Calix[4]arenes*; The Royal Society of Chemistry: Cambridge, UK, 1989. (g) Diamond, D.; McKervey, M. A. *Chem. Soc. Rev.* **1996**, *15*. (h) Ikeda, A.; Shinkai, S. *Chem. Rev.* **1997**, *97*, 1713. (i) Gutsche, C. D. *Calix[4]arenes Revisited*; The Royal Society of Chemistry: Cambridge, UK, 1998; pp 164–165. (j) Ludwig, R. In *Calixarenes 2001*; Kluwer Academic: Dordrecht, The Netherlands, 2001; pp 598–611. (k) Lee, D. H.; Lee, K. H.; Hong, J.-I. *Org. Lett.* **2001**, *3*, 5.

(2) For 4-nitrophenylazophenol-coupled calix[4]arenes as chromogenic ionophores, see: (a) Shimizu, H.; Iwamoto, K.; Fujimoto, K.; Shinkai, S. *Chem. Lett.* **1991**, 2147. (b) McCarrick, M.; Harris, S. J.; Diamond, D. *J. Mater. Chem.* **1994**, *4*, 217. (c) Gordon, J. L. M.; Böhmer, V.; Vogt, W. *Tetrahedron Lett.* **1995**, *36*, 2445. (d) Yamamoto, H.; Ueda, K.; Sandanayake, K. R. A. S.; Shinkai, S. *Chem. Lett.* **1995**, 497. (e) Chawla, H. M.; Srinivas, K. *J. Org. Chem.* **1996**, *61*, 8464. (f) Kim, N. Y.; Chang, S.-K. *J. Org. Chem.* **1998**, *63*, 2362. (g) Ma, Q.; Ma, H.; Su, M.; Wang, Z.; Nie, L.; Liang, S. *Anal. Chim. Acta* **2001**, *439*, 73. (h) Oueslati, F.; Dumazet-Bonnamour, I.; Lamartine, R. *Tetrahedron Lett.* **2001**, *42*, 8177. (i) Halouani, H.; Dumazet-Bonnamour, I.; Lamartine, R. *Tetrahedron Lett.* **2002**, *43*, 3785. (j) Kim, J. S.; Shon, O. J.; Lee, J. K.; Lee, S. H.; Kim, J. Y.; Park, K.-M.; Lee, S.-S. *J. Org. Chem.* **2002**, *67*, 1372. (k) Kim, J. Y.; Kim, G.; Kim, C. R.; Lee, S. H.; Lee, J. H.; Kim, J. S. *J. Org. Chem.* **2003**, *68*, 1933. (l) Lee, S. H.; Kim, J. Y.; Ko, J.; Lee, J. Y.; Kim, J. S. *J. Org. Chem.* **2004**, *69*, 2902.

(3) (a) Talanova, G. G.; Hwang, H.-S.; Talanov, V. S.; Bartsch, R. A. *Chem. Commun.* **1998**, 419. (b) Talanova, G. G.; Hwang, H.-S.; Talanov, V. S.; Bartsch, R. A. *Chem. Commun.* **1998**, 1329.

(4) (a) Cobben, P. L. H. M.; Egberink, R. J. M.; Bomer, J. G.; Bergveld, P.; Verboom, W.; Reinhoudt, D. N. *J. Am. Chem. Soc.* **1992**, *114*, 10573. (b) van der Veen, N. J.; Egberink, R. J. M.; Engbersen, J. F. J.; van Veggel, F. J. C. M.; Reinhoudt, D. N. *Chem. Commun.* **1999**, 681. (c) van der Veen, N. J.; Rozniecka, E.; Woldering, L. A.; Chudy, M.; Huskens, J.; van Veggel, F. J. C. M.; Reinhoudt, D. N. *Chem. Eur. J.* **2001**, *7*, 4878.

(5) (a) Shu, C.-M.; Lin, W.-L.; Lee, G.-H.; Peng, S.-M.; Chung, W.-S. *J. Chin. Chem. Soc.* **2000**, *47*, 173. (b) Compound **2** was prepared in a five-step synthetic sequence, see: Ho, Z.-C.; Ku, M.-C.; Shu, C.-M.; Lin, L.-G. *Tetrahedron* **1996**, *52*, 13189. (c) Shu, C.-M.; Yuan, T.-S.; Ku, M.-C.; Ho, Z.-C.; Liu, W.-C.; Tang, F.-S.; Lin, L.-G. *Tetrahedron*, **1996**, *52*, 9805. (d) Yeh, M.; Tang, F.; Chen, S.; Liu, W.; Lin, L.-G. *J. Org. Chem.* **1994**, *59*, 754.

(6) For a procedure of hydroboration, see: Moran, J. K.; Georgiev, E. M.; Yordanov, A. T.; Mague, J. T.; Roundhill, D. M. *J. Org. Chem.* **1994**, *59*, 5990.

(7) For lower rim esterification of calix[4]arenes, see: (a) Chen, X.; Wai, C. M.; Fisher, D. R. U.S. Patent 6,075,130, 2000. (b) Casnati, A.; Ting, Y.; Berti, D.; Fabbri, M.; Pochini, A.; Ungaro, R.; Sciotto, D.; Lombardo, G. G. *Tetrahedron* **1993**, *49*, 9815. (c) Chang, S.-K.; Cho, I. *J. Chem. Soc., Perkin Trans. 1* **1986**, 211. (d) McKervey, M. A.; Seward, E. M.; Ferguson, G.; Ruhl, B.; Harris, S. J. *J. Chem. Soc., Chem. Commun.* **1985**, 388. (e) Arduini, A.; Pochini, A.; Reverberi, S.; Ungaro, R.; Andreotti, G. D.; Ugozzoli, F. *Tetrahedron* **1986**, *42*, 2089. (f) Arimura, T.; Kubota, M.; Matsuda, T.; Manabe, O.; Shinkai, S. *Bull. Chem. Soc. Jpn.* **1989**, *62*, 1674.

(8) For some of the rare examples reporting complexation between azo groups on calix[4]arenes and transition metal ions see: (a) Nomura, E.; Taniguchi, H.; Otsuji, Y. *Bull. Chem. Soc. Jpn.* **1993**, *66*, 3797. (b) Nomura, E.; Taniguchi, H.; Tamura, S. *Chem. Lett.* **1989**, 1125. (c) References 2g and 2h. (d) Deligöz, H.; Erdem, E. *Solvent Ext. Ion Exch.* **1997**, *15*, 811.

(9) (a) Kishimoto, S.; Kitahara, S.; Manabe, O.; Hiyama, H. *J. Org. Chem.* **1978**, *43*, 3882. (b) Shinkai, S.; Araki, K.; Shibata, J.; Tsugawa, D.; Manabe, O. *Chem. Lett.* **1989**, 931. (c) Joshi, H.; Kamounah, F. S.; van der Zwan, G.; Gooijer, C.; Antonov, L. *J. Chem. Soc., Perkin Trans. 2* **2001**, 2303.





FIGURE 1. X-ray crystallographic structures of calix[4]arenes (a) **3a** and (b) **3c**.


SCHEME 1. Synthetic Pathways for Chromoionophores **3a–c and **4a–c**^a**



^a Reagents and conditions: (i) *p*-X-aniline/acetone, NaNO₂/4 M HCl, pyridine, 0 °C, 1 h.

3a–c, **4a–c**, and **5a,b**, their UV/vis spectra in CH₃CN and in CHCl₃ were inspected first. As expected, there were bathochromic shifts in λ_{\max} when piperidine (10 μ M) was added to the azocalix[4]arenes **3a–c**, **4a–c**, and **5a,b** (10–20 μ M in chloroform, Figure 2a–c); however, these changes in spectra were fully reversible, as the addition of HCl to these solutions reversed the color change. It should be noted that the shifts in λ_{\max} were small for **3a,b**, **4a,b**, and **5a,b** up to 40 mM of added NaOH, whereas large shifts in λ_{\max} were observed for the *p*-nitrophenyl azocalixarenes **3c–4c** when the same amount of NaOH was added. This is due to the ease in

SCHEME 2. Synthetic Pathways for Chromoionophores **5a,b and **6b,c**^a**



^a Reagents and conditions: (i) (1) 4 equiv of BH₃·THF, THF, rt, 1.5 h; (2) NaOH, H₂O₂, rt, 1.5 h. (ii) NaH (20 equiv), BrCH₂CO₂Et (20 equiv), THF, 50 °C, 8 h.

deprotonation of the lower rim hydroxy groups in **3c–4c** by a base, which favors the quinone forms. Lower rim etherification of these azocalixarenes should block this tautomerism pathway, and small change in spectra was expected. Indeed, the UV/vis spectrum stays almost the same when piperidine (10 μ M) was added to the solution of **6b** (Figure 2d).

We next investigated the affinity of these upper rim allyl- and azo-coupled calix[4]arenes (**3a–c**, **4a–c**, **5a,b**, and **6b,c**) for a series of transition metal ions (Cr³⁺, Ni²⁺, Cu²⁺, Cd²⁺, Hg²⁺, and Pb²⁺) and group I (Li⁺ and Na⁺) and group II (Ca²⁺ and Ba²⁺) metal ions in a methanol–chloroform (v/v = 1/399) cosolvent. Table 1 summarizes the bathochromic shifts of azocalix[4]arenes **3a–c**, **4a–c**, **5a,b**, and **6b,c** after adding 5 equiv of metal perchlorates. Only minor changes in UV/vis spectra were seen when 5 equiv of group I and II metal ions were added to **3b** (Figure 3a), whereas marked hyperchromic shifts




FIGURE 2. UV/vis spectra of azocalix[4]arenes (10 μM) in chloroform (solid line) and in the presence of 10 μM of piperidine (dashed line) (a) **3a**, (b) **3b**, (c) **3c**, and (d) **6b**.

TABLE 1. Bathochromic Shifts ($\Delta\lambda_{\text{max}}/\text{nm}$) of Azocalix[4]arenes **3a–c**, **4a–c**, **5a,b**, and **6b,c** by the Addition of 50 μM Metal Perchlorates in a Methanol–Chloroform ($v/v = 1/399$) Cosolvent^a

compd	λ_{max} , nm	$\Delta\lambda_{\text{max}}/\text{nm}^b$ after adding metal perchlorates									
		Li^+	Na^+	Ca^{2+}	Ba^{2+}	Cr^{3+}	Ni^{2+}	Cu^{2+}	Cd^{2+}	Hg_2^{2+}	Pb^{2+}
3a	348	143 ^c	143 ^c	142 ^c	133	136	127	132	129	132	129
3b	359	159	159	161	161	162	164	162	161	161	162
3c	377	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc
4a	349	nc	nc	141 ^c	141 ^c	134 ^c	140 ^c	140 ^c	128	131	128
4b	359	159	159	160	161	161	162	159	160	162	161
4c	377	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc
5a	347	130 ^c	130	130	130	135	128	133	130	138	130
5b	359	159	159	160	161	162	161	160	161	160	161
6b	359	-5 ^d	-6 ^d	-2	0	-3	-1	-1	0	-2	-2
6c	370	-22	-27	-17	-15	-17	-18	-15	-4	-8	-2

^a Concentration for **3a–c** and **5a,b** was 10 μM and that for **4a–c** was 20 μM , where nc denotes no detectable change in λ_{max} upon metal ion complexation. ^b $\Delta\lambda_{\text{max}} = \lambda_{\text{complex}} - \lambda_{\text{max}}(\text{free host})$. ^c Absorption at this λ_{complex} was below 0.02. ^d Besides the blue shift of the 359 nm band, there is a weak absorption band around 450–525 nm. ^e An apparent peak at λ_{max} around 485–495 nm was observed.

were observed for transition metal ions, especially for Hg^{2+} , Cr^{3+} , and Cu^{2+} ions (Figure 3b). It is noteworthy that 10 μM piperidine alone gave very little color change in **3b**; however, the light yellowish solution of **3b** turned into a bright red color upon complexation with $\text{Hg}(\text{ClO}_4)_2$, which is clearly visible to the naked eye (Figure 4). Surprisingly, no detectable changes in absorption spectra were observed for the very popular 4-(4-nitrophenyl)-azophenol calix[4]arenes **3c** and **4c** when 10 different metal ions were added in excess separately (see Table 1). These observations imply that there is a subtle balance between metal complexation-induced release of protons from the azophenols to the quinone-hydrazone tautomer.^{2,8,9} Moreover, the para substituent of the phenyl azo group must have a strong influence on the

tautomerism of azo/hydrazone, thus showing very diverse λ_{max} shifts. A picture of metal-ion- or base-induced tautomerism of the para-substituted phenyl azophenol is shown in Scheme 3.

Upon interacting with $\text{Hg}(\text{ClO}_4)_2$, the allyl- and azo-based chromoionophore **3b** in methanol–chloroform ($v/v = 1/399$) solution experienced a marked bathochromic shift in its λ_{max} as shown in Figure 5a. The absorption maximum at 359 nm gradually decreased in intensity with the formation of a new absorption band at ca. 520 nm ($\Delta\lambda_{\text{max}} = 161$ nm). Three isosbestic points at 263, 285, and 410 nm can be easily identified when difference absorption spectra analyses¹⁰ were used for the titration spectra of **3b** by $\text{Hg}(\text{ClO}_4)_2$ (Figure 5b). The spectra features in Figure 5 are consistent with a 1:1 binding




FIGURE 3. UV/vis spectra of **3b** (10 μM) before and after adding a 50 μM concentration of various metal perchlorates in a methanol–chloroform ($v/v = 1/399$) cosolvent: (a) Li^+ , Na^+ , Ca^{2+} , and Ba^{2+} ions; (b) Hg^{2+} , Cr^{3+} , Cu^{2+} , Ni^{2+} , and Pb^{2+} ions. The absorption band at 359 nm decreased in intensity with the formation of a new band at ca. 520 nm.





FIGURE 4. Colors of (a) **3b** (10 μM) in a methanol–chloroform ($v/v = 1/399$) cosolvent, (b) **3b** after adding 10 μM piperidine, and (c) **3b** after adding 5 equiv of $\text{Hg}(\text{ClO}_4)_2$.

SCHEME 3. Tautomerism of the Azophenol and Quinone-hydrazone



ratio between calix[4]arene **3b** and Hg^{2+} ion. Further support of the 1:1 binding ratio comes from a Job's plot experiment,¹¹ where the absorptions of the complex at 520 nm were plotted against molar fractions of **3b** under the conditions of an invariant total concentration. As a result, the concentration of **3b**· Hg^{2+} complex approached a maximum when the molar fraction of $[\mathbf{3b}]/([\mathbf{3b}] + [\text{Hg}^{2+}])$ was about 0.5 (see Figure 6).

(10) (a) Difference absorption spectra analysis is a popular method used in biophysical and biochemical fields, see: Lyles, M. B.; Cameron, I. L. *Biophys. Chem.* **2002**, *96*, 53. (b) Three isosbestic points were reported in alizarin-modified β -CD: Aoyagi, T.; Nakamura, A.; Ikeda, H.; Ikeda, T.; Mihara, H.; Ueno, A. *Anal. Chem.* **1997**, *69*, 659.

(11) Connors, K. A. *Binding Constants*, Wiley: New York, 1987.




FIGURE 5. (a) Changes in the UV/vis spectra of **3b** (10 μM) upon titration by $\text{Hg}(\text{ClO}_4)_2$ in a methanol–chloroform ($v/v = 1/399$) cosolvent, where the concentration of $\text{Hg}(\text{ClO}_4)_2$ varies from 1 to 14 μM , and (b) the difference absorption spectra of a.

Since very few changes in UV/vis spectra were found when azocalix[4]arenes **3a–c**, **4a–c**, and **5a,b** were treated with excess group I and II metal ions, their association constants were not determined. Among the eight upper rim azo-modified calix[4]arenes, only **3b**, **4b**, and **5b** showed marked bathochromic shifts upon complexation with transition metal ions (Cr^{3+} , Ni^{2+} , Cu^{2+} , and Hg^{2+}); therefore, their 1:1 association constants were determined on the basis of Benesi–Hilderbrand plots (see




FIGURE 6. Job's plot of a 1:1 complex of **3b** and Hg^{2+} ion, where the absorption at 520 nm was plotted against the mole fraction of **3b** at an invariant total concentration of 2×10^{-5} M in a MeOH/ CHCl_3 ($v/v = 1/19$) cosolvent.




FIGURE 7. Benesi-Hilderbrand plot of **3b** with $\text{Hg}(\text{ClO}_4)_2$.

TABLE 2. Association Constants (M^{-1}) in Log Scale ($\log K_a$) Based on UV/vis Titrations of Ionophores **3a,b**, **4a,b**, **5a,b**, and **6b,c** with Transition Metal Ion Perchlorates in a Methanol-Chloroform ($v/v = 1/399$) Cosolvent at 25 °C

compd	$\log K_a$, (M^{-1})			
	Cr^{3+}	Ni^{2+}	Cu^{2+}	Hg^{2+}
3a	a	a	a	5.17
3b	4.85	4.82	5.25	5.37
4a	a	a	a	4.70
4b	4.57	a	a	4.80
5a	a	a	a	4.73
5b	4.98	4.92	4.94	5.07
6b	3.48	a	a	4.27
6c	a	a	a	a
	$\text{Li}^+ = 4.07^b$	$\text{Na}^+ = 4.37^b$		

^a Association constant was not determined because the absorption differences of the complex were too small to be evaluated.

^b Association constants for group I metal ions were measurable from UV/vis titrations.

Experimental Section).¹² A typical plot for **3b** and $\text{Hg}(\text{ClO}_4)_2$ is shown in Figure 7, and data for all association constants (K_a in log scale) are summarized in Table 2. It is noteworthy that most of the K_a s for phenyl azophenols **3a–5a** and *para*-nitrophenyl azophenols **3c–4c** were not determined because their metal-ion-induced absorption changes were mostly too small to be evaluated except for Hg^{2+} ion. In contrast, the *para*-methoxyphenyl azophenol

series (**3b**, **4b**, and **5b**) showed very high sensitivity and selectivity toward transition metal ions, especially Hg^{2+} ion. The relative binding strengths of these chromogenic ionophores toward Hg^{2+} ion were determined to be **3b** > **3a** > **5b** > **4b** > **5a** ~ **4a** > **6b** \gg **3c–6c**. Furthermore, the selectivity trends of **3b** toward cations were determined to be Hg^{2+} > Cu^{2+} > Cr^{3+} > Ni^{2+} \gg (Pb^{2+} , Cd^{2+} , Ca^{2+} , Ba^{2+} , Na^+ , and Li^+) as shown in Figure 3 and Table 2.

The hypsochromic shift of λ_{max} (359 nm) of **6c** by 22 and 27 nm when binding with Li^+ and Na^+ , respectively, implies that **6c** uses the lower rim tetra ether-esters to bind metal ions (Figures S-8 and S-9, Supporting Information), which is consistent with those reported by Kim for lower rim crown-ether-modified azocalix[4]arenes.^{2k,l} When complexed with metal ion, the oxygen atoms of the lower rim ether-ester units of **6c** are positively polarized; therefore, the excited states are more strongly destabilized by cations than the ground states,^{1a,13} thus causing blue shifts of the λ_{max} band. Had the lower rim ether-esters been the only binding sites, one would have observed similar chromogenic behaviors for the homologue azocalix[4]arenes **6b** and **6c**, because their only structural difference is in the upper rim: *para*-methoxyphenyl for **6b** and *para*-nitrophenyl for **6c**. However, a quite different UV band splitting was observed when **6b** complexed with metal ions, where both a small hypsochromic shift and a large bathochromic shift in λ_{max} appeared concomitantly (Table 1). Furthermore, UV/vis titration of **6b** by Hg^{2+} showed two λ_{max} around 479 and 492 nm, which implies that Hg^{2+} may be entrapped in two different binding sites (see Figure S-6, Supporting Information). All these results imply that *para*-methoxyphenyl azophenol is capable of binding metal ions through its quinone-hydrazone tautomer; therefore, marked bathochromic shifts are observed.

To gain insight into the structures of complex formation of these chromogenic ionophores, we also carried out ^1H NMR titration experiments. Figure 8 shows the ^1H NMR spectra of **3b** (5 mM) in methanol- d_4 / CDCl_3 ($v/v = 1:3$) solution in the presence of different amounts of Hg^{2+} ion. Three important features were observed from the ^1H NMR spectra: (1) Large variation in chemical shifts is experienced by protons on the *para*-methoxyphenyl and azophenol groups; that is, protons H_f and H_a on both ends of the azo groups move 0.10 ppm upfield to H_f' and H_a' upon complexation with Hg^{2+} (see inset of Figure 8). (2) Signal intensities of protons on the allyl groups (namely, H_c , H_d , and H_e) decreased dramatically (to less than 30% of its original) in the presence of 2.2 equiv of Hg^{2+} ion. (3) The vaguely seen broad singlets of the methylene bridges around δ 3.5–4.5 in free ligand **3b** were shifted upfield to δ 2–4 upon complexation with Hg^{2+} and became sharp and complicated.

It is conceivable that these changes in chemical shifts and signal intensities are due to coordination of the Hg^{2+} ion to the upper rim azo groups (in its hydrazone form) and that strong dipolar interactions occur between the Hg^{2+} ion and the nearby allyl groups. This binding mode of the complex not only explains the changes in chemical shifts but also explains why the allyl signals diminish

(12) Benesi; H. A.; Hilderbrand, J. H. *J. Am. Chem. Soc.* **1949**, *71*, 2703.

(13) Alternatively, metal ions may stabilize the ground states more than the excited states, thus causing a hypsochromic shift of **6c**.





FIGURE 8. ^1H NMR spectra of compound **3b** (5.0 mM) in a methanol- d_4 /CDCl $_3$ (1:3) solution in the presence of different amounts of $\text{Hg}(\text{ClO}_4)_2$: (a) 0, (b) 5.56 mM (1.1 equiv), and (c) 11.1 mM (2.2 equiv). The inset on the right is an expanded region of δ 6.9 to 8.0, where H_f and H_a were shifted upfield when 2.2 equiv of Hg^{2+} was added.

when 2 equiv of Hg^{2+} ion was added (Figure 8c) since it is well-known that spin-lattice relaxation time (T_1) is greatly shortened by the existence of a nearby paramagnetic species such as an Hg^{2+} ion.¹⁴ The role of *p*-methoxy in the arylazo group is vital here because it helps to donate electrons toward the azo group through resonance, which reinforces the hydrazone that is formed through deprotonation of the azophenol to quinone-hydrazone (structure **A** in Scheme 3).

Inspecting the data reported in Table 2, one may readily find that calix[4]arenes with two upper rim azo groups on a face-to-face positions (e.g., **3a,b** and **5a,b**) give higher association constants toward transition metal ions than those with only one upper rim azo and three allyl groups (e.g., **4a,b**). Furthermore, upper rim allyl groups do not seem to have been involved in the binding of metal ions; they behave like bystanders because no change in chemical shifts was observed upon metal ion complexation. On the basis of all the information available for the metal complexes of these upper rim azo-

(14) For references about the influence of T_1 by paramagnetic species, see: (a) Lambert, B.; Jacques, V.; Desreux, J. F. In *Calixarenes for Separations*; Lumetta, G. J., Rogers, R. D., Gopalan, A. S., Eds.; American Chemical Society: Washington, DC, 2000; Chapter 13. (b) Wu, G.; Jiang, W.; Lamb, J. D.; Bradshaw, J. S.; Izatt, R. M. *J. Am. Chem. Soc.* **1991**, *113*, 6538. (c) Gutsche, C. D.; Iqbal, M.; Alam, I. *J. Am. Chem. Soc.* **1987**, *109*, 4314.

CHART 1. Possible Structure of the **3b**· Hg^{2+} Complex



coupled calix[4]arenes, we propose a possible structure for the **3b**· Hg^{2+} complex as shown in Chart 1. It is noteworthy that these metal complexes are stabilized by two pairs of intramolecular hydrogen-bonding interactions between undissociated phenolic OH groups (of *p*-allylphenol) and neighboring dissociated phenolate anions. Further support of the **3b**· Hg^{2+} complex structure comes from ^1H NMR and IR spectra (see Figures S-17 and S-18, Supporting Information).¹⁵ NMR spectra of free and Hg^{2+} -complexed **3b** using CD $_3$ OH/CDCl $_3$ (3/1) as a

(15) We thank one of the referees for suggestions of these NMR and IR experiments, which are very informative with respect to the structures of the complex.

cosolvent were compared with those in Figure 8 using CD₃OD/CDCl₃ (3/1) as a cosolvent. In CD₃OH/CDCl₃ cosolvent, the phenolic protons of free host **3b** showed a broad peak at δ 10.4 that disappeared when complexed with 2 equiv of Hg²⁺ ion. On the other hand, two new signals around δ 7.82 and 7.70 flanking the Ha and Ha' signals appeared, which are quite likely to be signals of hydrazone protons and the intramolecular phenol protons (inset c, Figure S-17, Supporting Information). IR spectra of the **3b**·Hg²⁺ complex were also consistent with the azophenol to quinone–hydrazone transformation (Figure S-18, Supporting Information).

Summary and Conclusions

All of these sensors are based on the well-known azobenzene structure, where one of the aromatic rings is an integrated part of the ion receptor. In the series of **3b–5b**, where 4-(4-methoxyphenyl)azophenol was used, high sensitivity of these chromoionophores for Hg²⁺ ion were found in a methanol–chloroform (v/v = 1/399) cosolvent. The Hg²⁺ ion detection gives rise to a large bathochromic shift in the absorption spectrum (from light yellow to bright red), which is clearly visible to the naked eye. The structure in Chart 1 provides a reasonable metal ion binding mode for the 4-phenyl- and 4-(4-methoxyphenyl)azophenol calix[4]arenes **3a,b** and **5a,b**, respectively. No shifts in absorption spectra were observed for the very popular 4-(4-nitrophenyl)azophenol calix[4]arenes **3c–4c** when 10 different metal ions were added in excess. The lack of binding of **3c–4c** may be explained by the predominant form of structure **B** in Scheme 3, in which the strongly electron-withdrawing nitro group prevents the formation of the hydrazone form; therefore, poor binding toward metal cations was observed. To the best of our knowledge, these upper rim allyl-, hydroxypropyl-, and azo-coupled calix[4]arenes are rare examples⁸ of such highly sensitive Hg²⁺ ion sensors based on azobenzene-calix[4]arene alone without further modification on the lower rims.

Experimental Section

The syntheses of 5,11-bisallylcalix[4]arene **1**,^{5c} 5,11,17-triallylcalix[4]arene **2**,^{5b} and 5,17-bisallyl-11,23-bis(phenylazo)-calix[4]arenes **3a–c**^{5c} have been reported by Lin et al., and we have followed their procedures for the preparation.

General Procedure for the Synthesis of Allylaryl-azo-25,26,27,28-tetrahydroxy calix[4]arenes 3a–c^{5c} and 4a–c. To an ice cold solution of 4.0 mmol of para-substituted aniline in 5 mL of 4 N HCl was added a solution of 0.41 g (5.94 mmol) of NaNO₂ in 5 mL of acetone, and the mixture was stirred for 30 s. The combined solution was then added to another ice cold solution of 0.50 g (1.0 mmol) of *p*-diallyl- or *p*-triallylcalix[4]arenes (**1** or **2**) in 10 mL of pyridine to produce a colored solution. The reaction mixture was stirred for 1 h at 0 °C and then treated with 50 mL of 4 N HCl to give a colored precipitate. The solid residue was purified by column chromatography with hexane and chloroform as an eluent and gave the corresponding bisarylozo or monoarylozo products in 61–91% yields.

Data for 5,11,17-Triallyl-23-phenylazo-25,26,27,28-tetrahydroxy calix[4]arene, 4a. The solid was eluted with hexane/chloroform (v/v = 2/1) and gave 0.55 g (91%) of an orange powder, mp 293–297 °C (dec): ¹H NMR (300 MHz, CDCl₃) δ 10.21 (s, 4H, OH), 7.81 (d, 2H, *J* = 7.1 Hz), 7.68 (s, 2H), 7.51–7.42 (m, 3H), 6.95–6.85 (m, 6H), 5.92–5.82 (m, 3H),

5.08–4.99 (m, 6H), 4.35–4.15 (bs, 4H), 3.70–3.40 (bd, 4H), 3.21–3.16 (m, 6H); ¹³C NMR (75.4 MHz, CDCl₃) δ 152.8 (C_q), 151.9 (C_q), 147.9 (C_q), 146.9 (C_q), 146.8 (C_q), 137.4 (CH), 133.8 (C_q), 133.5 (C_q), 130.3 (CH), 129.2 (CH), 129.0 (CH), 128.9 (CH), 128.3 (C_q), 128.0 (C_q), 127.5 (C_q), 124.0 (CH), 122.4 (CH), 115.7 (CH₂), 115.6 (CH₂), 39.3 (CH₂), 31.8 (CH₂), 31.7 (CH₂); FAB-MS *m/z* 649 (M + H⁺); HRMS *m/z* calcd for C₄₃H₄₀N₂O₄ 648.2990, found 648.3033. Anal. Calcd for C₄₃H₄₀N₂O₄: C, 79.59; H, 6.22; N, 4.32. Found: C, 79.38; H, 6.32; N, 4.53.

Data for 5,11,17-Triallyl-23-(*p*-methoxyphenyl)azo-25,26,27,28-tetrahydroxy calix[4]arene, 4b. The solid was eluted with hexane/chloroform (v/v = 2/1) and gave 0.49 g (78%) of a yellow powder, mp 260–262 °C (dec): ¹H NMR (300 MHz, CDCl₃) δ 10.20 (s, 4H, OH), 7.85–7.81 (m, 2H), 7.63 (s, 2H), 7.00–6.94 (m, 4H), 6.87–6.84 (m, 4H), 5.92–5.80 (m, 3H), 5.08–4.99 (m, 6H), 4.35–4.15 (bs, 4H), 3.88 (s, 3H), 3.70–3.40 (bd, 4H), 3.21–3.16 (m, 6H); ¹³C NMR (CDCl₃) δ 161.6 (C_q), 151.3 (C_q), 147.5 (C_q), 146.9 (C_q), 137.5 (CH), 133.8 (C_q), 133.6 (C_q), 129.2 (CH), 129.0 (CH), 128.8 (CH), 128.3 (C_q), 128.0 (C_q), 127.6 (C_q), 124.3 (CH), 123.6 (CH), 115.6 (CH₂), 114.1 (CH), 55.5 (CH₃), 39.3 (CH₂), 31.8 (CH₂), 31.7 (CH₂); FAB-MS *m/z* 679 (M + H⁺); HRMS *m/z* calcd for C₄₄H₄₂N₂O₅ 678.3096, found 678.3110. Anal. Calcd for C₄₄H₄₂N₂O₅: C, 77.84; H, 6.24; N, 4.13. Found: C, 77.76; H, 6.40; N, 4.22.

Data for 5,11,17-Triallyl-23-(*p*-nitrophenyl)azo-25,26,27,28-tetrahydroxy calix[4]arene, 4c. The solid was eluted with hexane/chloroform (v/v = 2/1) and gave 0.47 g (73%) of a deep red powder, mp 256–259 °C (dec): ¹H NMR (300 MHz, CDCl₃) δ 10.22 (s, 4H, OH), 8.37–8.33 (m, 2H), 7.95–7.92 (m, 2H), 7.73 (s, 2H), 6.95–6.85 (m, 6H), 5.92–5.80 (m, 3H), 5.08–5.00 (m, 6H), 4.40–4.20 (bs, 4H), 3.70–3.40 (bd, 4H), 3.22–3.16 (m, 6H); ¹³C NMR (CDCl₃) δ 156.0 (C_q), 153.4 (C_q), 148.2 (C_q), 147.3 (C_q), 146.9 (C_q), 146.8 (C_q), 137.4 (CH), 133.9 (C_q), 133.7 (C_q), 129.3 (CH), 129.3 (CH), 129.1 (CH), 129.0 (CH), 128.4 (C_q), 127.9 (C_q), 127.2 (C_q), 124.7 (CH), 123.1 (CH), 115.7 (CH₂), 115.6 (CH₂), 39.3 (CH₂), 31.8 (CH₂), 31.7 (CH₂); FAB-MS *m/z* 694 (M + H⁺); HRMS *m/z* calcd for C₄₃H₃₉N₃O₆ 693.2841, found 693.2823. Anal. Calcd for C₄₃H₃₉N₃O₆: C, 74.43; H, 5.67; N, 6.06. Found: C, 74.04; H, 5.68; N, 6.25.

General Procedure for the Synthesis of 5,17-Bis(3-hydroxypropyl)-11,23 bisarylozocalix[4]arenes 5a,b.⁶ To a solution of **3a** (60 mg, 0.080 mmol) in tetrahydrofuran (7 mL) was added BH₃·THF (0.30–0.33 mL of 1 M solution) dropwise over a period of 20 min. The solution was stirred for 1 h at room temperature, and then water (2 mL) was added slowly to destroy excess BH₃. The pH was raised by addition of 3 mL of sodium hydroxide (3 M aqueous solution), and then H₂O₂ (3 mL of a 30% solution) was added dropwise over 10 min. The solution was heated to 30–50 °C for 15 min and then stirred at room temperature for 1 h. The solution was then extracted with ethyl acetate (3 × 30 mL), and the combined extracts were washed with water (2 × 30 mL) and brine (50 mL) and then dried over magnesium sulfate. The solution was removed, and the residue was purified by column chromatography with hexane and ethyl acetate as an eluent and gave the corresponding 3-hydroxypropyl-arylazocalix[4]arenes **5a** and **5b** in 39 and 58% yields, respectively.

Data for 5,17-Bis(3-hydroxypropyl)-11,23-bis(phenyl)azocalix[4]arenes 5a. The solid was eluted with hexane/ethyl acetate (v/v = 1/1) and gave 24 mg (39%) of an orange powder, mp 263–267 °C (dec): ¹H NMR (300 MHz, CDCl₃) δ 10.22 (bs, 4H, OH), 7.82–7.79 (m, 4H), 7.68 (s, 4H), 7.50–7.41 (m, 6H), 7.05–7.01 (m, 4H), 4.40–4.28 (bd, 4H), 3.72–3.55 (m, 8H), 2.57–2.51 (m, 4H), 1.86–1.76 (m, 4H); ¹³C NMR (CDCl₃) δ 152.7 (C_q), 151.8 (C_q), 147.5 (C_q), 146.6 (C_q), 135.9 (C_q), 130.4 (CH), 129.3 (CH), 129.0 (CH), 128.8 (C_q), 127.6 (CH), 124.0 (CH), 122.5 (CH), 62.0 (CH₂), 34.0 (CH₂), 31.8 (CH₂), 31.0 (CH₂); FAB-MS *m/z* 750 (M + H⁺); HRMS *m/z* calcd for C₄₆H₄₄N₄O₆ 748.3263, found 748.3242.

Data for 5,17-Bis(3-hydroxypropyl)-11,23-bis(*p*-methoxyphenyl)azocalix[4]arenes 5b. The solid was eluted with hexane/ethyl acetate (v/v = 1/1) and gave 36 mg (58%) of

a yellow powder, mp 267–271 °C (dec): ^1H NMR (300 MHz, CDCl_3) δ 10.20 (bs, 4H, OH), 7.83–7.79 (m, 4H), 7.63 (s, 4H), 7.02–6.94 (m, 8H), 4.37–4.23 (bd, 4H), 3.86 (s, 6H), 3.70–3.53 (m, 8H), 2.56–2.50 (m, 4H), 1.83–1.75 (m, 4H); ^{13}C NMR (CDCl_3) δ 161.6 (C_q), 151.2 (C_q), 147.5 (C_q), 147.0 (C_q), 146.6 (C_q), 135.8 (C_q), 129.3 (CH), 128.8 (C_q), 128.0 (C_q), 124.3 (CH), 123.6 (CH), 114.1 (CH), 62.0 (CH_2), 55.5 (CH_3), 34.0 (CH_2), 31.8 (CH_2), 31.0 (CH_2); FAB-MS m/z 809 ($\text{M} + \text{H}^+$); HRMS m/z calcd for $\text{C}_{48}\text{H}_{48}\text{N}_4\text{O}_8$ 808.3474, found 808.3369.

General Procedures for the Lower Rim Esterification of the Azo **3b and **3c** to Give Compounds **6b** and **6c**.** The method is adapted from a procedure reported by Chang^{7c} et al. for lower rim esterification of calix[4]arene. Compound **3b** or **3c** (30 mg, 0.037–0.042 mmol) was dissolved in dry THF (4 mL), and then ethyl bromoacetate (0.1 mL, 0.75–0.84 mmol, 20 equiv) and NaH (20 mg, 0.75–0.84 mmol, 20 equiv) were added under nitrogen. The resulting mixture was stirred at 50 °C for 9 h. After the mixture was cooled to room temperature, the solvent was removed under reduced pressure, and the residue was redissolved in chloroform and washed with water. The chloroform solution was dried over MgSO_4 and evaporated to give the solid crude product. Column chromatography on silica gel eluting with hexane/ethyl acetate ($v/v = 4/1$) gave the desired tetraesters **6b** and **6c** in 74 and 36% yields, respectively.

Data for 5,17-Diallyl-11,23-bis(*p*-methoxyphenyl)azo-25,26,27,28-tetrakis((ethoxycarbonyl)methoxy)calix[4]-arene **6b.** A yellow solid of 32 mg (74%) was obtained, mp 150–152 °C: ^1H NMR (300 MHz, CDCl_3) δ 7.78–7.74 (m, 4H), 7.44 (s, 4H), 6.93–6.88 (m, 4H), 6.45 (s, 4H), 5.79–5.64 (m, 2H), 4.94–4.81 (m, 12H), 4.66 (s, 4H), 4.29–4.17 (m, 8H), 3.90–3.87 (m, 6H), 3.35 (d, 4H, $J = 13.6$ Hz), 2.97 (d, 4H, $J = 6.4$ Hz), 1.35–1.27 (m, 12H); ^{13}C NMR (75.4 MHz, CDCl_3) δ 170.1 (C_q), 169.9 (C_q), 161.5 (C_q), 158.6 (C_q), 153.7 (C_q), 148.2 (C_q), 147.0 (C_q), 137.8 (CH), 135.9 (C_q), 134.5 (C_q), 133.2 (C_q), 128.7 (CH), 124.3 (CH), 123.2 (CH), 115.1 (CH_2), 114.0 (CH), 71.6 (CH_2), 71.2 (CH_2), 60.6 (CH_2), 60.5 (CH_2), 55.4 (CH_3), 39.3 (CH_2), 31.5 (CH_2), 14.2 (CH_3), 14.2 (CH_3); FAB-MS m/z 1118 ($\text{M} + 2\text{H}^+$), 1140 ($\text{M} + \text{H}^+ + \text{Na}^+$); HRMS ($\text{M} + \text{H}^+$) m/z calcd for $\text{C}_{64}\text{H}_{69}\text{N}_4\text{O}_{14}$ 1117.4813, found 1117.4801.

Data for 5,17-Diallyl-11,23-bis(*p*-nitrophenyl)azo-25,26,27,28-tetrakis((ethoxycarbonyl)methoxy)calix[4]-arene **6c.** A red solid of 15 mg (36%) was obtained, mp 180–183 °C: ^1H NMR (300 MHz, CDCl_3) δ 8.17–8.13 (m, 4H), 7.76–7.71 (m, 4H), 7.35 (s, 4H), 6.60 (s, 4H), 5.86–5.77 (m, 2H), 4.97–4.85 (m, 12H), 4.73 (s, 4H), 4.28–4.18 (m, 8H), 3.36 (d, 4H, $J = 13.7$ Hz), 3.09 (d, 4H, $J = 6.2$ Hz), 1.34–1.23 (m, 12H); ^{13}C NMR (CDCl_3) δ 170.0 (C_q), 169.7 (C_q), 159.9 (C_q), 155.6

(C_q), 154.1 (C_q), 148.1 (C_q), 148.0 (C_q), 137.8 (CH), 135.9 (C_q), 134.6 (C_q), 133.7 (C_q), 129.1 (CH), 124.4 (CH), 124.0 (CH), 122.9 (CH), 115.2 (CH_2), 71.5 (CH_2), 71.3 (CH_2), 39.2 (CH_2), 31.6 (CH_2), 14.2 (CH_3), 14.6 (CH_3); FAB-MS m/z 1148 ($\text{M} + 2\text{H}^+$), 1170 ($\text{M} + \text{H} + \text{Na}^+$); HRMS m/z calcd for $\text{C}_{62}\text{H}_{62}\text{N}_6\text{O}_{16}$ 1146.4225, found 1146.4252.

General Procedures for UV/vis Experiments. Because of the poor solubility of metal perchlorates in chloroform, all of the UV/vis experiments reported in this work were carried out in a methanol–chloroform ($v/v = 1/399$) cosolvent unless otherwise specified. UV/vis spectra were recorded on an HP-8453 spectrophotometer with a diode array detector, and the resolution was set at 1 nm. The complexes were mixed and kept stirring for at least 20 min before measurements. During the measurements, the temperature of the quartz sample cell and the chamber was kept at 25 °C.

Stability Constants of Metal Complexes. The stability constants K_a of a 1:1 complex of **3a–c**, **4a–c**, and **5a,b** with four transition metal ions (Cr^{3+} , Ni^{2+} , Cu^{2+} , and Hg^{2+}) were determined from the following Benesi–Hilderbrand equation:¹²

$$1/\Delta A = 1/\Delta A_{\text{sat}} + 1/(\Delta A_{\text{sat}} K_a [\text{guest}]) \quad (1)$$

where ΔA is the absorption difference between the apparent metal complex and the free ionophores and ΔA_{sat} is the absorption difference at saturation. The association constant values K_a were evaluated graphically by plotting $1/\Delta A$ against $1/[\text{metal ion}]$. In this study, the concentration of the ionophores was kept constant (10 μM for **3a–c** and **5a,b** and 20 μM for **4a–c**), and the concentration of metal ions was kept in the range $[\text{metal ion}]/[\text{ionophores}] = 0.1\text{--}2.0$. All plots gave a straight line, and the K_a values were obtained from the slopes and intercepts of this line (See Supporting Information).

Acknowledgment. We thank the National Science Council of the Republic of China for financial support (Grants NSC 90-2113-M-009-017 and 91-2113-M-009-011).

Supporting Information Available: IR spectra, UV/vis titration spectra, Job's plots, and Benesi–Hilderbrand plots of **3b** and **6b** with various transition metal ions and **6c** with Li^+ and Na^+ ions; partial X-ray crystallography data for compounds **3a** and **3c**; ^1H NMR spectra of compounds **3a–c**, **5a,b**, and **6b,c**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO047880A